ELSEVIER

Journal of Organometallic Chemistry 485 (1994) C27–C28

Journal of Organometallic Chemistry

Author Index of Volume 485

Abboud, K.A., 37
Adatia, T., 191
Akita, M., C14
Akiyama, T., 31
Alt, H.G., 11
Amor, J.I., 153
Ang, H.-G., C10
Ang, S.-G., C10
Arena, C.G., 115
Arif, A.M., 25

Battistuzzi, R., 257
Bell, W., 201
Blanco, L., 63
Blaschette, A., 179
Blum, J., 209
Bochkarev, L.N., 101
Boennemann, H., 31
Boncella, J.M., 37
Brunner, H., 237
Bruno, G., 115
Burton, N.C., 153

Cecchi, P., 45
Chakrabarti, P., C1
Chernega, A., C18
Churakov, A.V., 173
Colapietro, M., 45
Cuenca, T., 153

Denisov, V.R., 219
Djerourou, A., 63
Doak, G.O., 1
Dolgushin, F.M., 219, 253
Drommi, D., 115

El Amouri, H., 79
Erabi, T., 127
Ernst, R.D., 25
Ershova, V.A., 69

Fan, W.-Q., 19
Faraone, F., 115
Fierro, R., 11
Fischer, R., 227
Freedman, L.D., 1
Fröhlich, H.-O., 227
Fukin, G.K., 101

Gavioli, G., 257
Gehart, G., 237
Gioia Lobbia, G., 45
Glidewell, C., 201
Gómez-Sal, P., 153
Goren, N., 209
Green, M.L.H., C18
Gruselle, M., 79

Hassler, K., 233
Hatano, H., 31
Hayase, S., 127
Hemling, H., 209
Herrman, G.S., 11
Herrmann, W.A., 243
Hill, C.M., 165
Hirakawa, H., C14
Hu, J., C6
Huang, X., C6
Huo, S.Q., 161

Igarashi, A., 31
Iguchi, Y., 31

Jaouen, G., 79
Jones, P.G., 179
Joshi, V.S., C1

Kaganovich, V.S., 109
Kajitani, M., 31
Kanzaki, M., 127
Karpov, M.G., 219
Kazennova, N.B., 173
Kerzina, Z.A., 109
Kolehmainen, E., 109
Kollegger, G., 233
Kondo, T., 55
Korvola, J., 109
Kühn, F.E., 243
Kuz'mina, L.G., 173
Kwik, W.-L., C10

Labella, L., C18
Laihia, K., 109
Lange, I., 179
Leblanc, J.-C., 237
Lemenovskii, D.A., 173
Li, X.-W., 141, 149
Lindeman, S.V., 257
Lorberth, J., 141, 149

Maksakov, V.A., 69
Mansour, B., 123
Mason, S.J., 165
Massa, W., 141, 149
Massey, S.T., 123
Matsumoto, S., 31
McAndrew, D., 201
McCarthy, P.J., 191
McElwee-White, L., 123
McGlinchey, M.J., 79
Mitsudo, T., 55
Moise, C., 237
Molosnova, N.E., 101
Moro-oka, Y., C14
Morrison, C.A., 201
Murphy, V.J., 165

Nagy, A., 215
Nandi, M., C1
Nicolò, F., 115
Nikol'skii, A.B., 219
Nuber, B., 237

Ogura, H., 127
O'Hare, D., 165
Osintseva, S.V., 253

Pályi, G., 257
Pettinari, C., 45
Pifferi, A., 45
Pozdniakova, Yu.A., 257

Ramm, M., 135
Rausch, M.D., 11
Roesky, P.W., 243
Royo, P., 153
Ruan, M.-D., 19
Rufanov, K.A., 173
Rybin, L.V., 253
Rybinskaya, M.I., 109, 253

Salter, I.D., 191
Santi, P., 257
Sarkar, A., C1
Sarraje, I., 149
Sathe, K.M., C1
Satô, G.P., 31
Schäfer, M., 85
Schumann, H., 209

Selent, D., 135
Shchegolikhina, O.I., 257
Shen, F., C6
Shimizu, K., 31
Shin, S.-H., 149
Spagna, R., 45
Starova, G.L., 219
Struchkov, Y.T., 101
Struchkov, Yu.T., 219, 253
Sugimori, A., 31
Szepes, L., 215

Tanaka, M., C14
Thöne, C., 185
Trakarnpruk, W., 25
Tunik, S.P., 219

Ugo, R., 257

Vahrenkamp, H., 185
Vaissermann, J., 79
Vass, G., 215
Vaughan, W.M., 37
Vizi-Orosz, A., 257
Volpato, F., 237

Wachter, J., 237
Wada, M., 127
Wang, S., C6
Watanabe, Y., 55
Watkin, D.J., 165
Werner, H., 85
Wocadlo, S., 141, 149
Wolf, J., 85
Wu, Y.J., 161
Wyrwa, R., 227

Xie, M., C6

Yanovsky, A.I., 101, 219, 253

Zakaria, C.M., 201
Zakharov, L.N., 101
Zanathy, L., 215
Zhang, Q., C10
Zhang, S.-W., 55
Zhao, H.-R., 19
Zhdanov, A.A., 257
Zhou, X.-J., 19
Zhu, Y., 161
Zucchi, C., 257

ELSEVIER

Journal of Organometallic Chemistry 485 (1994) C29–C36

Journal of Organometallic Chemistry

Subject Index of Volume 485

Acetato complexes
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85
Acetylide(alkynyl)
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-}C{\equiv}C\text{-}Fp)Ru_3(\mu\text{-}H)(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
Alcohol
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
Alkinyl(hydrido) complexes
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85
Alkoxides
Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[ZrCp_2(OC_6F_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153
Alkyl halide
First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19
Alkylidenes
Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $TpMo(CH_2C(Me)_2Ph)(NAr)(O)$ and $[TpMo(NAr)(O)]_2O$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37
Alkyne complexes
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85
Allyl
Severe distortion of π-allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π-cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
Amine
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
Ansa-bridging
Ansa-[(2,2-bis-η-cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18
Arene
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^1H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Aryl
Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141
Aryl bismuth compounds
Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141
Aurated clusters
Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185
Azides
Reaction of $(CO)_5W(THF)$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123

Binuclear complexes
Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115
2,2-bis-η-cyclopentadienyl(propane)
Ansa-[(2,2-bis-η-cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18

Bismuth
Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141
Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, (^{1}h-$C_5H_5)_3Bi$" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149
Borate
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
Boron
He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215
Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of $K(HB(3,4,5\text{-}Me_3Pz)_3]$ and $[HB(3,4,5\text{-}Me_3Pz)_3\ SnMeCl_2]$ (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
Bridging ligand
Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115
Bulky ligands
Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Cage
Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-Cl})\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
Carbanions
Synthesis and structure of triphenylmethyl derivatives of ytterbium $[Yb(THF)_6]^{2+}[Ph_3C]_2^-$ and $[(THF)_4Yb(\mu\text{-Cl})_2Yb(THF)_4]^{2+}\ [Ph_3Cl]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101
Carbonyl
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^{1}H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Carboxylic acid synthesis
Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von $[Li(TMEDA)]_2\overline{Ni(CH_2CH_2CH_2CH_2C(O)O)}_2$ und seine Umsetzung mit CH_3I und einem C_{22}-Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227
Catalysis
Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethanamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethanaminium]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209
Catalyst
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
C–C-bond formation
Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von $[Li(TMEDA)]_2\overline{Ni(CH_2CH_2CH_2CH_2C(O)O)}_2$ und seine Umsetzung mit CH_3I und einem C_{22}-Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227
Charge-transfer
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron-(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
Chromium
Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^{1}H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Cluster
Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-Cl})\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-C}\equiv\text{C–Fp})Ru_3(\mu\text{-H})(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10
Cobalt
Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185
Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-Cl})\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^{1}H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^{1}H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Reactions of diacetylene ligands with trinuclear clusters. I. Reactions of 2,4-hexadiyne-1,6-diol and its dicobalthexacarbonyl derivatives with $H_2Os_3(CO)_{10}$ (M.G. Karpov, S.P. Tunik, V.R.

Denisov, G.L. Starova, A.B. Nikol'skii, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 219

Cone angles

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}\ \{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Coupling

Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-}C{\equiv}C\text{–}Fp)Ru_3(\mu\text{-}H)(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Crystal structure

Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Polysulfonylamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-*N*)dimethylzinn-(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179

Severe distortion of π-allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π-cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetonato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron-(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}\ \{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Cyclohexane

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

Cyclometallated derivative

Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Cyclopentadienyl

Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[ZrCp_2(OC_6F_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, (^{1}h-$C_5H_5)_3Bi$" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetonato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6

Cyclopolyphosphine

Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Cyclosiloxane

Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-}Cl)\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Dicarbide

Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-}C{\equiv}C\text{–}Fp)Ru_3(\mu\text{-}H)(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Dichlorosilanes

Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63

Diketones

Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201

Dimesylamide

Polysulfonylamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-*N*)dimethylzinn-(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179

2,6-Dimethoxyphenyl

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Diselenides

First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19

Disulfide ligand

Comparative studies in the $Cp_2M(\eta^2\text{-}S_2)H$ series (Cp = t-BuC_5H_4, C_5Me_4Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Ferrocene

Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201

Ferrocenes
Synthesis and characterization of $Fe(1\text{-}Me_3Si\text{-}3\text{-}CH_3C_5H_5)_2$, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25
Ferrocenylimine
Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
Functionalised polysilanes
Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane $MeSi(SiMe_2X)_3$ und Hexasilane $(Me_2XSi)_2MeSiSiMe(SiMe_2X)_2$, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233

Group 4
Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[ZrCp_2(OC_6F_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153
Group 11
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}\{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
Group 15
Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, $(^1h\text{-}C_5H_5)_3Bi$" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Hafnium
He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215
Heterobimetallics
Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-}Cl)\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
Holmium
Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetonato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6
Hunsdiecker reaction
Surprisingly facile decomposition of the dication $[(C_5Me_5)Ir(MeO\text{–}C_6H_4\text{–}CH_2CO_2\overline{NCOCH_2CH_2CO})]^{2+}$: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79
Hydride ligand
Comparative studies in the $Cp_2M(\eta^2\text{-}S_2)H$ series (Cp = t-BuC_5H_4, C_5Me_4Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237
Hydrogenation
Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethanamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethanaminium]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209
Hydrogen bonding
Molekül- und Kristallstruktur von μ–Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{–}OH)Rh(COD\text{-}1,5)]_2$ mit $[\eta^5\text{–}C_5(CH_3)_5]_2Ti(IV)(CH_3)_2$ und $(\eta^5\text{–}C_5H_5)_2Ti(II)[P(CH_3)_3]_2$ (D. Selent und M. Ramm), 135
Hydrolysis
Polysulfonylamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-*N*)dimethylzinn-(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179
Hydroxide
Polysulfonylamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-*N*)dimethylzinn-(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179
Hydroxylamine
Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Iridium
Surprisingly facile decomposition of the dication $[(C_5Me_5)Ir(MeO\text{–}C_6H_4\text{–}CH_2CO_2\overline{NCOCH_2CH_2CO})]^{2+}$: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79
Iron
Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185
Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-}C{\equiv}C\text{–}Fp)Ru_3(\mu\text{-}H)(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron-(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
Isomerization
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Ketenylidene-Fe_3-cluster
Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185
Ketenylidene-Fe_2Co-cluster
Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185

Lanthanide complexes
Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetonato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6
Synthesis and structure of triphenylmethyl derivatives of ytterbium $[Yb(THF)_6]^{2+}[Ph_3C]_2^-$ and $[(THF)_4Yb(\mu\text{-}Cl)_2Yb(THF)_4]^{2+}$ $[Ph_3Cl]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101
Lithium enolates
Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63

Magnetic susceptibility
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
Mercury
Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
Metallocenes
Comparative studies in the $Cp_2M(\eta^2\text{-}S_2)H$ series (Cp = t-BuC_5H_4, C_5Me_4Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237
Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11
Molybdenum
Ansa-[(2,2-bis-η-cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18
Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69
Severe distortion of π-allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π-cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1
Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $TpMo(CH_2C(Me)_2Ph)(NAr)(O)$ and $[TpMo(NAr)(O)]_2O$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37

2-Naphthoyltrifluoroacetonato
Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetonato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6
Nickel
Bimetallic siloxane cluster of higher valent transition metals: $Na\{[\eta^6\text{-cyclo-}(PhSiO_2)_6]_2Co_2Ni_4(\mu_6\text{-}Cl)\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $CpSn(CH_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-$(CH_3)_3Si$–(3-Ph_3C)–C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173
Nickelaoxacycloheptanones
Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von $[Li(TMEDA)]_2\dot{N}i(CH_2CH_2CH_2CH_2C(O)\dot{O})_2$ und seine Umsetzung mit CH_3I und einem C_{22}-Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227
Nitrenes
Reaction of $(CO)_5W(THF)$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123
^{17}O-NMR spectroscopy
Mehrfachbindungen zwischen Übergangsmetallen und Hauptgruppenelementen. CXXXIV. ^{17}O-NMR-Spektroskopie an Organorhenium(VII)-oxiden (W.A. Herrmann, F.E. Kühn und P.W. Roesky), 243
Nuclear magnetic resonance
Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^1H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Severe distortion of π-allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π-cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Open ferrocenes
Synthesis and characterization of $Fe(1\text{-}Me_3Si\text{-}3\text{-}CH_3C_5H_5)_2$, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25
Organorhenium(VII) oxides
Mehrfachbindungen zwischen Übergangsmetallen und Hauptgruppenelementen. CXXXIV. ^{17}O-NMR-Spektroskopie an Organorhenium(VII)-oxiden (W.A. Herrmann, F.E. Kühn und P.W. Roesky), 243
Osmium
Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69
Reactions of diacetylene ligands with trinuclear clusters. I. Reactions of 2,4-hexadiyne-1,6-diol and its dicobalthexacarbonyl derivatives with $H_2Os_3(CO)_{10}$ (M.G. Karpov, S.P. Tunik, V.R. Denisov, G.L. Starova, A.B. Nikol'skii, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 219
Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Palladium
Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
Pentadienyl complexes
Synthesis and characterization of $Fe(1\text{-}Me_3Si\text{-}3\text{-}CH_3C_5H_5)_2$, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25

Phosphine
Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethanamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethanaminium]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209
Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^{1}H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
Phosphine ligands
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}\ \{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
Phosphonium
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
Phosphorus
Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10
Photoelectron spectroscopy
He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215
Platinum
Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115
Properties
Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane $MeSi(SiMe_2X)_3$ und Hexasilane $(Me_2XSi)_2MeSiSiMe(SiMe_2X)_2$, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233
Pyrazde
Severe distortion of π-allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π-cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1
Pyrazolylborate
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
Pyrazolylborates
Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of $K(HB(3,4,5\text{-}Me_3Pz)_3]$ and $[HB(3,4,5\text{-}Me_3Pz)_3\ SnMeCl_2]$ (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45
Reactivity
Comparative studies in the $Cp_2M(\eta^2\text{-}S_2)H$ series (Cp = t-BuC_5H_4, C_5Me_4Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237
Red modification
Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, (^{1}h-$C_5H_5)_3Bi$" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149
Rhodium
Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethanamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethanaminium]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209
Molekül- und Kristallstruktur von μ–Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{–}OH)Rh(COD\text{-}1,5)]_2$ mit $[\eta^5\text{–}C_5(CH_3)_5]_2Ti(IV)(CH_3)_2$ und $(\eta^5\text{–}C_5H_5)_2Ti(II)[P(CH_3)_3]_2$ (D. Selent und M. Ramm), 135
Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85
Ruthenium
Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^{1}H, ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109
Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115
Reactions of dodecacarbonyltriruthenium with enones: η^3-coordinated dihydropyranyl rings (L.V. Rybin, S.V. Osintseva, M.I. Rybinskaya, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 253
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-}C{\equiv}C\text{–}Fp)Ru_3(\mu\text{-}H)(CO_9)]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-}C_2H_x)M_y$ cluster compounds ($x = 0, 1$; $y = 7, 8$), $CpFeRu_6(\mu_5\text{-}C_2)(\mu_5\text{-}C_2H)(CO)_{16}$, $Cp_2Fe_2Ru_5(\mu_5\text{-}C_2)_2(CO)_{17}$ and $Cp_2Fe_2Ru_6(\mu_6\text{-}C_2)_2(CO)_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β-keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}\ \{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
Selenium
First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl dise-

lenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19

Selenoamides

First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19

Selenourea

First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19

3-Silaglutarates

Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63

Silicon

Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $CpSn(CH_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-$(CH_3)_3Si$–(3-Ph_3C)–C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173

Silver

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[Ag_2Ru_4(\mu_3\text{-}H)_2(CO)_{12}$ $\{P(C_6H_4Me\text{-}2)_3\}_2]$. An investigation of the steric properties of the $P(C_6H_4Me\text{-}2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Solubility

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Sulfur

Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Synthesis

Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane $MeSi(SiMe_2X)_3$ und Hexasilane $(Me_2XSi)_2MeSiSiMe(SiMe_2X)_2$, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233

Tantalum

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215

TCNQ

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron-(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165

Thiolate

Formation constants of some phosphine and phosphite adducts of (η^5-cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H, ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Tin

Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of $K(HB(3,4,5\text{-}Me_3Pz)_3]$ and $[HB(3,4,5\text{-}Me_3Pz)_3\ SnMeCl_2]$ (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45

Polysulfonylamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-*N*)dimethylzinn-(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179

Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $CpSn(CH_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-$(CH_3)_3Si$–(3-Ph_3C)–C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173

Titanium

Molekül- und Kristallstruktur von μ–Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{–}OH)Rh(COD\text{-}1,5)]_2$ mit $[\eta^5\text{–}C_5(CH_3)_5]_2Ti(IV)(CH_3)_2$ und $(\eta^5\text{–}C_5H_5)_2Ti(II)[P(CH_3)_3]_2$ (D. Selent und M. Ramm), 135

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[ZrCp_2(OC_6F_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11

Triphenylmethyl derivatives

Synthesis and structure of triphenylmethyl derivatives of ytterbium $[Yb(THF)_6]^{2+}[Ph_3C]_2^-$ and $[(THF)_4Yb(\mu\text{-}Cl)_2Yb(THF)_4]^{2+}$ $[Ph_3Cl]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

2,4,6-Triphenylphenyl

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Tris-cyclopentadienyl bismuth

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, $(^1h\text{-}C_5H_5)_3Bi$" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Trispyrazolylborates

Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $TpMo(CH_2C(Me)_2Ph)(NAr)(O)$ and $[TpMo(NAr)(O)]_2O$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37

Tungsten

Ansa-[(2,2-bis-η-cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Reaction of $(CO)_5W(THF)$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123

Vinylidene complexes

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkinyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

X-ray

Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of

K(HB(3,4,5-$Me_3Pz)_3$] and [HB(3,4,5-$Me_3Pz)_3$ $SnMeCl_2$] (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45

X-ray analysis

Molekül- und Kristallstruktur von μ–Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von [(μ–OH)Rh(COD-1,5)]$_2$ mit [η^5–$C_5(CH_3)_5$]$_2$Ti(IV)$(CH_3)_2$ und (η^5–$C_5H_5)_2$Ti(II)[$P(CH_3)_3$]$_2$ (D. Selent und M. Ramm), 135

X-ray diffraction

Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethanamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethanaminium]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of [$ZrCp_2(OC_6F_5)_2$] (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, (^{1}h-$C_5H_5)_3$Bi" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Synthesis and structure of triphenylmethyl derivatives of ytterbium [$Yb(THF)_6]^{2+}[Ph_3C]_2^-$ and [$(THF)_4Yb(\mu$-$Cl)_2Yb$-$(THF)_4]^{2+}$ $[Ph_3Cl]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

X-ray structure

Aurated clusters derived from (μ_3-ketenylidene)-Fe_3 and -Fe_2Co clusters (C. Thöne and H. Vahrenkamp), 185

Comparative studies in the $Cp_2M(\eta^2$-$S_2)H$ series (Cp = t-BuC_5H_4, C_5Me_4Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Mononuclear(η^6-arene)ruthenium(II) and (η^5-pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)–platinum(II) and ruthenium(II)–rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Surprisingly facile decomposition of the dication [(C_5Me_5)Ir-$(MeO$–C_6H_4–$CH_2CO_2\overline{NCOCH_2CH_2CO})]^{2+}$: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79

Ytterbium

Synthesis and structure of triphenylmethyl derivatives of ytterbium [$Yb(THF)_6]^{2+}[Ph_3C]_2^-$ and [$(THF)_4Yb(\mu$-$Cl)_2Yb$-$(THF)_4]^{2+}$ $[Ph_3Cl]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

Zirconium

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215

Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of [$ZrCp_2(OC_6F_5)_2$] (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Book Reviews

Catalysis of Organic Reactions, John R. Kosak and Thomas A. Johnson (eds.) (J.R. Hanson), C22

Leaving No Stone Unturned: Pathways in Organometallic Chemistry F. Gordon A. Stone (*Profiles, Pathway, and Dreams: Autobiographies of Eminent Chemists*, Jeffrey I. Seeman, Series Editor) (M.F. Lappert), C22

Molybdenum Enzymes, Cofactors, and Model Systems, E.I. Stiefel, D. Coucouvanis and W.E. Newton (Eds.) (P.G.J. Leigh), C22